

COVID-19 Burden and Control in 2020: A Multi-Metric Analysis Across Five Countries

Jacob P. Cherian ¹, , Jubilant J. Kizhakkethottam ²

1. *Research Scholar, APJ Abdul Kalam Technological University, Thiruvananthapuram, IND*

2. *Computer Science and Engineering, Saintgits College of Engineering, Kottayam, IND*

Received: September 04, 2025 | Review began: November 05, 2025 | Review ended: June 10, 2026 | Published: June 15, 2026

© Copyright 2026

This is an open access article distributed under the terms of the Creative Commons Attribution License CC-BY 4.0., which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

We analyse the COVID-19 transmission, severity, and public health response in five countries: India, Brazil, Italy, the United Kingdom, and the United States. The analysis includes several key indicators: the effective reproduction number (R_t), confirmed cases and deaths per million population, case fatality rate, and a government policy stringency index. A derived mortality ratio is also included to assess severity relative to case incidence. Data are sourced from the Our World in Data repository and standardised to allow for temporal comparison across countries. The results show that Brazil and the United Kingdom experienced extended periods of uncontrolled transmission, with R_t remaining above 1 for much of the year. Italy maintained shorter intervals over this threshold because policy enforcement was earlier and more frequent. The case fatality rate showed significant variation, with the United Kingdom having the highest mean rate and India having the lowest. Many nations did not show a consistent correlation between transmission and policy rigor. Although the indicators do not take into consideration all structural and behavioral factors, they do offer a comparative view of national trajectories. The findings provide insight into the relationship between disease outcomes, timing of interventions, and transmission dynamics.

Categories: Health Informatics, Computational Social Science, Data Analysis

Keywords: covid 19, stringency index, reproduction rate, public health, case fatality rate

Introduction

Background and global context

When Coronavirus Disease 2019 (COVID-19) first appeared in late 2019, it quickly became a global health issue. The virus was able to spread even when people did not have symptoms or before they showed symptoms, which is why the outbreak started in Wuhan, China, and quickly spread to other countries. By March 2020, most countries had reported cases and were putting emergency measures in place to stop the spread. More and more patients, problems with the supply chain, not enough testing capacity, and not enough staff are all putting more and more stress on health systems.

The pandemic not only had direct health effects, but it also showed that public health readiness, data infrastructure, and governance were not strong enough. Governments had to act quickly, often based on changing clinical and epidemiological data, because science was unclear. In many places, lockdowns, travel bans, and school closures became common. However, the timing and effectiveness of these interventions were completely unique. Because of this, 2020 was a very important time to learn about how countries dealt with the first waves of the pandemic and what happened next. Looking at these answers gives us important information about how diseases spread and how well policies work.

How to cite this article:

Cherian J P, Kizhakkethottam J J (June 15, 2026) COVID-19 Burden and Control in 2020: A Multi-Metric Analysis Across Five Countries. Cureus J Comput Sci 3 : es44389-025-00079-3. DOI <https://doi.org/10.7759/s44389-025-00079-3>

Epidemiological framework

The effective reproduction number (R_t) is a common way to talk about how quickly COVID-19 spreads. It shows how many secondary infections a single case caused on average at a certain time. Mathematically, R_t represents the transmission potential at time t , derived from case growth rates and disease serial interval. If the values stay above 1, it means that the outbreak is getting worse. If the values stay below 1, it means that the epidemic is getting better. It has been very important to keep an eye on R_t to guide public health actions [1,2]. Several studies have shown that starting and sticking with non-pharmaceutical interventions (NPIs) early on can help bring R_t below the epidemic threshold [3,4].

The case fatality rate (CFR) and other severity measures have helped us learn even more about the national-level burden. The CFR changes depending on things such as the population age structure, how easy it is to obtain medical care, and how well the testing system works. CFR is advantageous but should be read carefully because reports may take longer to come in, case definitions may not be the same across countries, and case detection completeness varies substantially [5,6]. The CFR depends critically on testing rates, time delays in reporting, age structure of the population, and ascertainment bias [5,6]. You can also use deaths per million people or derived mortality ratios to compare how much disease is affecting people in different countries with different population sizes and ways of reporting.

Several attempts have created composite indices to evaluate and compare government policy responses. The Oxford COVID-19 Government Response Tracker is one of the most popular tools. The Oxford COVID-19 Government Response Tracker provides a daily stringency index that measures the adherence and strictness of various NPIs [7]. Some studies that looked at the link between stringency and epidemic trends found a link with R_t and case incidence, while others found that the effects were less strong once community spread was established [8,9]. This suggests that policy stringency is one of many factors influencing epidemic control, and implementation quality and population adherence may matter more than policy existence alone. Many studies have looked at country-level dynamics on their own, but fewer have used empirical data to apply the same metrics across multiple high-burden countries. A lot of the early writing was about either predictive modeling or short-term studies in specific areas [10,11].

Research gap and study objectives

This study attempts to fill this gap by using the same set of epidemiological and policy indicators in five countries that had big early outbreaks: India, Brazil, Italy, the United Kingdom, and the United States. These countries have a wide range of public health capacities, governance structures, and economic resources. While comparative analyses of cross-country metrics exist in the literature, this study provides a systematic integration of multiple standardized indicators over the complete 2020 calendar year, emphasizing the disconnect between policy stringency and actual epidemic control in some national contexts and examining how policy implementation quality and timing differ from policy existence.

This study looks at COVID-19 control outcomes during the first year of the pandemic from a comparative perspective by looking at R_t , CFR, confirmed case and death rates, policy stringency, and a derived mortality ratio. Earlier works have also assessed transmission control measures, early outbreak forecasting, and long-term epidemic trajectories, which provide useful background for interpreting cross-country patterns [12-15].

Materials And Methods

Data sources and indicators

The Our World in Data COVID-19 repository [15], which gathers daily, country-level data from official reports from national and international health agencies, provided all datasets used for this study. Data were extracted on June 15, 2025, ensuring complete 2020 coverage. The following indicators were chosen:

Primary Epidemiological Indicators

How to cite this article:

1. Rt: The Rt represents the mean number of secondary infections generated by a single infected individual at time t. Data sources: Our World in Data aggregates estimates from published models for each country. Mathematically, Rt is derived from case growth rates and disease serial interval. Values > 1 indicate exponential growth; values < 1 indicate epidemic decline.
2. Confirmed Cases per Million and Deaths per Million: These numbers are adjusted for population size by using cumulative values, which makes it possible to compare countries. Formulae: Cases per million = (Cumulative confirmed cases × 1,000,000)/Population; Deaths per million = (Cumulative deaths × 1,000,000)/Population.
3. CFR: $CFR(t) = [Deaths\ at\ time\ t / Cases\ at\ time\ t] \times 100\%$. This is the percentage of confirmed cases resulting in reported death. Testing rates, case definition consistency, reporting delays, and population age structure significantly influence the CFR.
4. Stringency Index: The Oxford COVID-19 Government Response Tracker uses this measure to show how strict policies are measured on a 0-100 scale, where 0 = no interventions and 100 = maximum stringency, including policy categories such as closing schools, banning travel, and running public information campaigns.
5. Derived Mortality Ratio: $MR(t) = Deaths\ per\ million / Cases\ per\ million$, ranging from 0 to 1. This metric attempts to account for incidence when comparing severity across countries. We note that this ratio inherits the same confounders as CFR.

Data processing and cleaning

Five countries were selected: India, Brazil, Italy, the United Kingdom, and the United States. These countries were chosen because they faced significant early waves and represent diverse health system capacities, governance structures, and demographic profiles. Data processing was carried out using Python 3.9.1 with the Pandas 1.3.0 and NumPy 1.21.0 libraries.

Processing steps:

1. Country name standardization before merging datasets to ensure proper alignment.
2. Rows with missing values in key indicators were excluded unless reporting continuity was confirmed (defined as ≥3 consecutive non-missing values).
3. Forward filling was applied only when reporting continuity was confirmed; no interpolation or extrapolation was done for periods with missing data to avoid introducing artificial trends.
4. CFRs and mortality ratios were recalculated from cumulative case and death counts to ensure internal consistency across countries.
5. Seven-day rolling averages were applied to Rt, confirmed cases, and deaths to minimize daily fluctuations, reporting irregularities, and weekend effects. Padding at series edges was applied using the forward/backward filling of boundary values.
6. The cleaned series were merged into a single panel indexed by date and country.

Experimental setup

This study covers the period from January 1, 2020 to December 31, 2020, which represents the first full year of the COVID-19 pandemic. Five countries were chosen: India, Brazil, Italy, the United Kingdom, and the United States. These countries were selected because they reported high case counts early and represent varied health system capacities and policy responses.

The analysis focused on five indicators: Rt, confirmed cases per million, deaths per million, CFR, mortality ratio, and the government stringency index. Data were taken from Our World in Data, which compiles daily figures from national and international health agencies.

How to cite this article:

Data preprocessing was carried out using Python 3.9.1 with Pandas 1.3.0 and NumPy 1.21.0. Rows with missing values were excluded unless reporting continuity was confirmed, in which case forward filling was applied. CFR and mortality ratio were recalculated from raw case and death counts to maintain consistency. Seven-day rolling averages were applied to Rt, case counts, and deaths to smooth daily variation and reporting artifacts.

The cleaned datasets were merged into a single panel indexed by date and country. Epidemic wave duration was measured by counting the number of days where Rt was greater than 1.0. All plots were prepared using Matplotlib (version 3.4.3) to visualize Rt trends, stringency levels, cases, deaths, and severity metrics. This setup allowed for a uniform and reproducible comparison of epidemic patterns across the five countries.

Results And Discussion

Limitations

Data Quality and Reporting Heterogeneity

Our analysis relies on country-level aggregate data characterized by substantial heterogeneity in reporting practices and surveillance capacity. Variations in testing infrastructure and case detection protocols across study countries introduce important measurement considerations. India experienced documented underreporting of mild and asymptomatic cases in 2020 owing to constrained testing infrastructure, whereas Brazil exhibited marked regional disparities in testing capacity between affluent states such as São Paulo and resource-limited northeastern regions. The United Kingdom implemented rapid testing protocols earlier than comparator nations (July 2020), potentially enhancing ascertainment of mild cases, while the United States demonstrated considerable state-level variation in testing policies alongside relatively comprehensive testing availability. Italy, among the earliest European nations to deploy mass testing, likely achieved more complete case ascertainment than other study settings. Death reporting similarly varied: India historically undercounted mortality in 2020, with excess mortality analyses suggesting substantial underestimation of true deaths; attribution criteria for COVID-19 mortality differed across jurisdictions (encompassing confirmed, probable, and suspected classifications); and reporting delays ranging from days to weeks rendered 2020 mortality counts incomplete at the time of analysis.

Consequently, CFR comparisons should be interpreted as reflecting within-country temporal trends rather than absolute cross-national rankings. India's comparatively low CFR likely reflects detection bias rather than genuine differences in disease severity, whereas the United Kingdom's elevated CFR may partially reflect more sensitive case detection combined with an older population structure. Mortality ratios inherit these same measurement constraints. Although Our World in Data applies standardization procedures to mitigate heterogeneity, residual variation persists; we therefore frame findings as relative burden trajectories rather than absolute severity metrics and advise caution in cross-country interpretation.

Structural and Demographic Confounders

Our quantitative framework does not fully account for structural heterogeneities that may influence observed associations. Healthcare system characteristics varied considerably: the United Kingdom and Italy operate publicly funded systems that experienced substantial capacity strain during the spring 2020 surge; the United States maintains a mixed public-private system with pronounced geographic disparities in intensive care unit capacity and resource allocation; Brazil's fragmented system exhibits marked inequality between affluent and underserved regions; and India faced limited ICU capacity in 2020, particularly outside major urban centers. Population demographics further modulate severity risk: the United Kingdom (median age approximately 40 years) and Italy (approximately 47 years) possess older age structures associated with elevated COVID-19 severity, whereas Brazil and India exhibit younger median ages (approximately 34 and 27 years, respectively), correlating with lower case fatality risk. Socioeconomic and governance factors also introduce unmeasured confounding: India's large informal economy (approximately 45% of the workforce in 2020) likely attenuated lockdown effectiveness despite high stringency indices; Brazil's political polarization undermined

How to cite this article:

adherence to centralized public health directives; the United Kingdom's dense public transportation networks may have amplified transmission independent of policy measures; and the United States experienced coordination challenges between federal and state authorities that created implementation gaps. While these factors are qualitatively acknowledged, our analysis could not quantitatively integrate them. Future work incorporating healthcare capacity indices, per capita testing rates, demographic standardization, and behavioral compliance metrics would strengthen causal inference.

Temporal Scope Limitation

This analysis is restricted to the calendar year 2020, a deliberate scope choice with both rationale and constraints. The 2020 focus addresses our specific research question concerning initial policy responses during the first full year of the pandemic, precedes the availability of vaccines (which would necessitate fundamentally different analytical frameworks), and maintains methodological consistency regarding case definitions and variant nomenclature. However, this temporal boundary precludes examination of policy evolution during 2021-2022, assessment of vaccination campaign impacts on transmission dynamics, and evaluation of how policy effectiveness may have shifted with the emergence of novel variants.

Rt Estimation Heterogeneity

Rt estimates utilized in this analysis are derived from Our World in Data's aggregation of multiple published methodological approaches. This introduces several limitations: different countries may employ distinct statistical methods to estimate Rt; aggregating across methodological frameworks introduces heterogeneity that is not fully characterized in secondary data sources; and Rt estimates are inherently dependent on the completeness of case detection, which varied substantially across countries and over time.

Causal Inference Limitations

Although our analysis identifies temporal associations between policy stringency indices and Rt trajectories, we cannot establish causal relationships. This limitation arises from the observational nature of the data, the presence of multiple unmeasured confounders (including voluntary behavioral adaptations, healthcare system capacity, demographic structure, and variant prevalence), and the potential for reverse causality (wherein elevated Rt values may prompt policy intensification rather than policies driving Rt reduction). Isolating the independent effect of policy measures from these intertwined factors requires methodological approaches beyond the scope of this descriptive analysis. Future studies employing instrumental variable strategies, difference-in-differences designs, or natural experiment frameworks would be better positioned to disentangle policy effects from confounding influences and strengthen causal inference regarding non-pharmaceutical intervention effectiveness.

Confirmed cases and deaths

The United States had the most confirmed cases per million people overall, followed by the United Kingdom and Brazil (Figure 1). The United Kingdom had the most deaths per million people (Figure 2). India had the fewest cases per person, even though it has a very large population.

How to cite this article:




FIGURE 1: Confirmed COVID-19 Cases Per Million

How to cite this article:

Cherian J P, Kizhakkethottam J J (June 15, 2026) COVID-19 Burden and Control in 2020: A Multi-Metric Analysis Across Five Countries. Cureus J Comput Sci 3 : es44389-025-00079-3. DOI <https://doi.org/10.7759/s44389-025-00079-3>




FIGURE 2: Confirmed COVID-19 Deaths Per Million

Temporal Analysis

There was a clear time gap between the peaks in cases and deaths in different countries, but the magnitude and implications differed. Italy experienced its peak case burden in April 2020, followed by a sharp decline and brief plateauing period (June-July) before a second wave in October-November. Brazil showed sustained high transmission from April through December, with multiple minor peaks. The United Kingdom exhibited a pronounced first peak in April and a more substantial second peak in December. The United States demonstrated multiple overlapping regional waves, with peaks in July and December. India's trajectory showed a delayed peak relative to Europe, with sustained but moderate transmission throughout the second half of 2020.

Explanatory Factors

The strength of the health system, the age composition of the population, and how complete the reporting of cases and deaths could all be reasons for these differences. However, we acknowledge that differences in testing capacity, case definitions, and reporting standards substantially confound direct cross-country severity comparisons.

CFR analysis

At the beginning of 2020, the United Kingdom had the highest CFR (Figure 3), with some weeks having more than 10%. This likely reflects the older age structure of the UK population (median age ~40 years) combined with healthcare system saturation during the spring 2020 peak and possibly incomplete case ascertainment in early months. Italy also had a high CFR during its first wave, reflecting similar demographic and healthcare capacity factors. Brazil's CFR stayed relatively stable, while India's and the United States's declined over time.

How to cite this article:




FIGURE 3: COVID-19 Case Fatality Rate

However, the United States may have been substantially affected by expanded testing revealing more mild cases over time, artificially lowering CFR. The fact that CFR is declining in most countries over time suggests that both case detection and treatment are improving, but the fact that there are still significant differences shows that healthcare systems are not equal and that demographic factors substantially influence outcomes.

Key Limitation

CFR is highly dependent on testing coverage, case definition consistency, reporting delays, age structure, and ascertainment bias. Cross-country CFR comparisons should be interpreted as relative trends within countries rather than absolute severity rankings. The United Kingdom's high CFR likely reflects both demographic factors (older population) and healthcare system stress, while India's low CFR likely reflects incomplete case detection in 2020 rather than intrinsically milder disease.

Stringency index and transmission control

All countries enacted stringent responses by late March. The UK (Figure 4), Italy (Figure 5), and Brazil (Figure 6) implemented early nationwide measures. United States (Figure 7) and India (Figure 8) exhibited greater variability, with subnational differences in policy enforcement.

How to cite this article:




FIGURE 4: United Kingdom - Rt vs Stringency Index

Rt, Effective Reproduction Rate




FIGURE 5: Italy - Rt vs Stringency Index

Rt, Effective Reproduction Rate




FIGURE 6: Brazil - Rt vs Stringency Index

Rt, Effective Reproduction Rate




FIGURE 7: United States - R_t vs Stringency Index

R_t , Effective Reproduction Rate




FIGURE 8: India - Rt vs Stringency Index

Rt, Effective Reproduction Rate

Mortality ratio analysis

The United Kingdom had the highest ratio at the beginning of the pandemic, surpassing 0.2 by April 2020 before progressively falling throughout the year (Figure 9). The same pattern, albeit at lower levels, was seen in Italy, which stabilized in the middle of 2020 at 0.15. Both extremes point to a severe initial wave that caused a significant number of recorded infections to result in deaths.

Demographic and Healthcare Context

The United Kingdom's and Italy's high initial mortality ratios reflect:

- (1) older age structures (median ages ~40 and ~47 years, respectively);
- (2) healthcare system saturation during spring 2020;
- (3) early waves occurring before treatment protocols improved.

In contrast, the ratios for Brazil and the United States were more intermediate, remaining between 0.05 and 0.1 and gradually decreasing by the end of 2020. India also had the lowest ratio, remaining consistently below 0.05. These patterns most likely stem from variations in the population [age structure], medical care [capacity], and case detection. We note that India's low ratio likely reflects underdetection of mild cases in 2020 due to limited testing infrastructure, rather than intrinsically lower disease severity.

How to cite this article:

Cherian J P, Kizhakkethottam J J (June 15, 2026) COVID-19 Burden and Control in 2020: A Multi-Metric Analysis Across Five Countries. *Cureus J Comput Sci* 3 : es44389-025-00079-3. DOI <https://doi.org/10.7759/s44389-025-00079-3>




FIGURE 9: COVID-19 Mortality Rate

Epidemic wave duration and transmission dynamics

The virus continued to spread in Brazil (Figure 10) and the United States for an extended period, with $R_t > 1$ for over 200 days. India had the fewest at 157 (Figure 11), even though total cases were high. Italy (Figure 12) managed to bring R_t down below 1.0 more quickly; Brazil had 233 such days, the United Kingdom (Figure 13) had 214. The United States (Figure 14) had a sharp decline in R_t by May 2020. R_t trends over time showed that the outbreak did not develop in the same way in every country (Figure 15).

Temporal Offset Explanation

This temporal variation reflects distinct epidemiological contexts:

- (1) Italy experienced the earliest wave (March-April), likely due to direct importations from China and rapid community spread in densely populated regions, followed by effective intervention;
- (2) The United Kingdom had delayed outbreak recognition (first UK deaths in mid-March) compared to Italy, resulting in later peak control;
- (3) The United States experienced multiple overlapping regional waves due to federal-state coordination gaps and geographic heterogeneity in intervention timing;
- (4) Brazil maintained sustained transmission due to documented federal-subnational policy disagreements;
- (5) India showed a delayed epidemic trajectory, possibly due to younger population demographics and the substantial informal economy reducing lockdown compliance despite high stringency.

Mortality Implications

How to cite this article:

Cherian J P, Kizhakkethottam J J (June 15, 2026) COVID-19 Burden and Control in 2020: A Multi-Metric Analysis Across Five Countries. *Cureus J Comput Sci* 3 : es44389-025-00079-3. DOI <https://doi.org/10.7759/s44389-025-00079-3>

Places with longer waves often ended up with more deaths too, consistent with transmission dynamics theory. The 76-day difference between Brazil (233 days) and Italy (157 days) with $R_t > 1$ corresponds approximately to 1-2 additional epidemic doubling periods, representing substantially increased cumulative transmission exposure.




FIGURE 10: Epidemic Waves for Brazil (with $R_t > 1$)

R_t , Effective Reproduction Rate




FIGURE 11: Epidemic Waves for India (with $R_t > 1$)

R_t , Effective Reproduction Rate




FIGURE 12: Epidemic Waves for Italy (with $R_t > 1$)

R_t , Effective Reproduction Rate




FIGURE 13: Epidemic Waves for United Kingdom (with $R_t > 1$)
 R_t , Effective Reproduction Rate




FIGURE 14: Epidemic Waves for United States (with $R_t > 1$)

R_t , Effective Reproduction Rate




FIGURE 15: Countries and Number of Days with $R_t > 1$

R_t , Effective Reproduction Rate

Conclusions

The analysis shows that among the five countries, Brazil and the United Kingdom had the longest sustained transmission, as indicated by R_t . Even though the outbreak was severe and early, Italy was better at keeping things under control for a longer period of time. There were several overlapping epidemic waves in the United States due to regional and geographic variations. Despite having one of the highest absolute case counts in the world, the epidemic trajectory of India was unique in that it had a delayed peak and shorter wave duration. The United Kingdom had the highest cumulative deaths per million and year-round high case fatality rates. India, on the other hand, continued to have the lowest fatality metrics on both derived and relative measures. This does not indicate lower disease severity in India but rather reflects incomplete case detection in 2020.

This analysis has provided a more complex understanding of epidemic and disease control by combining multiple indicators. This study gives valuable insights into how various nations, especially the severely hit ones, dealt with the first year of the pandemic by combining measures of transmission, severity, mortality, and policy response. Key finding: Policy stringency alone is an insufficient measure of epidemic control effectiveness; implementation quality, national coordination, timing of interventions, and population-level behavioral responses all substantially influence outcomes. As

How to cite this article:

Cherian J P, Kizhakkethottam J J (June 15, 2026) COVID-19 Burden and Control in 2020: A Multi-Metric Analysis Across Five Countries. Cureus J Comput Sci 3 : es44389-025-00079-3. DOI <https://doi.org/10.7759/s44389-025-00079-3>

future work, we can also incorporate behavioral and political determinants, including trust metrics, media landscape analyses, and compliance surveys to better understand how governance and public response mediate policy effectiveness.

Additional Information

Author Contributions

All authors have reviewed the final version to be published and agreed to be accountable for all aspects of the work.

Concept and design: Jacob P. Cherian, Jubilant J. Kizhakkethottam

Acquisition, analysis, or interpretation of data: Jacob P. Cherian, Jubilant J. Kizhakkethottam

Drafting of the manuscript: Jacob P. Cherian, Jubilant J. Kizhakkethottam

Critical review of the manuscript for important intellectual content: Jacob P. Cherian, Jubilant J. Kizhakkethottam

Supervision: Jubilant J. Kizhakkethottam

Disclosures

Human subjects: All authors have confirmed that this study did not involve human participants or tissue. **Animal subjects:** All authors have confirmed that this study did not involve animal subjects or tissue. **Conflicts of interest:** In compliance with the ICMJE uniform disclosure form, all authors declare the following: **Payment/services info:** All authors have declared that no financial support was received from any organization for the submitted work. **Financial relationships:** All authors have declared that they have no financial relationships at present or within the previous three years with any organizations that might have an interest in the submitted work. **Other relationships:** All authors have declared that there are no other relationships or activities that could appear to have influenced the submitted work.

Acknowledgements

This article was previously presented at the International Conference on NEXT GENERATION TECHNOLOGY IN SCIENCE AND ENGINEERING (ICNGTSE - 2025) at Pranveer Singh Institute of Technology, Kanpur (PSIT) on 22–23 August 2025.

References

1. Wu J, Leung K, Leung G: [Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study](#). The Lancet. 2020, 395:689-697. [10.1016/s0140-6736\(20\)30260-9](#)
2. Eubank S, Eckstrand I, Lewis B, Venkatramanan S, Marathe M, Barrett CL: [Commentary on Ferguson, et al., "Impact of Non-pharmaceutical Interventions \(NPIs\) to Reduce COVID-19 Mortality and Healthcare Demand"](#). Bulletin of Mathematical Biology. 2020, 82:52. [10.1007/s11538-020-00726-x](#)
3. Flaxman S, Mishra S, Gandy A, et al.: [Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe](#). Nature. 2020, 584:257-261. [10.1038/s41586-020-2405-7](#)
4. Hsiang S, Allen D, Annan-Phan S, et al.: [The effect of large-scale anti-contagion policies on the COVID-19 pandemic](#). Nature. 2020, 584:262-267. [10.1038/s41586-020-2404-8](#)
5. Verity R, Okell L, Dorigatti I, et al.: [Estimates of the severity of coronavirus disease 2019: a model-based analysis](#). The Lancet Infectious Diseases. 2020, 20:669-677. [10.1016/s1473-3099\(20\)30243-7](#)
6. Dong E, Du H, Gardner L: [An interactive web-based dashboard to track COVID-19 in real time](#). The Lancet Infectious Diseases. 2020, 20:533-534. [10.1016/s1473-3099\(20\)30120-1](#)
7. Hale T, Petherick A, Phillips T: [Variation in government responses to COVID-19 \(Version 15, BSG Working Paper Series, BSG-WP-2020/032\)](#). (2023). <https://www.bsg.ox.ac.uk/sites/default/files/2023-06/BSG-WP-2020-032-v15.pdf>.

How to cite this article:

Cherian J P, Kizhakkethottam J J (June 15, 2026) COVID-19 Burden and Control in 2020: A Multi-Metric Analysis Across Five Countries. Cureus J Comput Sci 3 : es44389-025-00079-3. DOI <https://doi.org/10.7759/s44389-025-00079-3>

8. Wang G, Zhang Y, Zhao J, Zhang J, Jiang F: [Mitigate the effects of home confinement on children during the COVID-19 outbreak](#). The Lancet. 2020, 395:945-947. [10.1016/s0140-6736\(20\)30547-x](#)
9. Zhang J, Litvinova M, Liang Y, et al.: [Changes in contact patterns shape the dynamics of the COVID-19 outbreak in China](#). Science. 2020, 368:1481-1486. [10.1126/science.abb8001](#)
10. Pei S, Kandula S, Shaman J: [Differential effects of intervention timing on COVID-19 spread in the United States](#). Science Advances. 2020, 6:eabd6370. [10.1126/sciadv.abd6370](#)
11. Gatto M, Bertuzzo E, Mari L, Miccoli S, Carraro L, Casagrandi R, Rinaldo A: [Spread and dynamics of the COVID-19 epidemic in Italy: Effects of emergency containment measures](#). Proceedings of the National Academy of Sciences. 2020, 117:10484-10491. [10.1073/pnas.2004978117](#)
12. Tian H, Liu Y, Li Y, et al.: [An investigation of transmission control measures during the first 50 days of the COVID-19 epidemic in China](#). Science. 2020, 368:638-642. [10.1126/science.abb6105](#)
13. Kissler SM, Tedijanto C, Goldstein E, Grad YH, Lipsitch M: [Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period](#). Science. 2020, 368:860-868. [10.1126/science.abb5793](#)
14. Tang B, Wang X, Li Q, Bragazzi NL, Tang S, Xiao Y, Wu J: [Estimation of the transmission risk of the 2019-nCoV and its implication for public health interventions](#). Journal of Clinical Medicine. 2020, 9:462. [10.3390/jcm9020462](#)
15. Mathieu E, Ritchie H, Rod  s-Guirao L, et al.: [COVID-19 pandemic](#). (2020). Accessed: June 15, 2025: <https://ourworldindata.org/coronavirus>.

How to cite this article: